

T-Wave Alternans and Autonomic Nervous System Activity During Orthostatic Stress after 5 Days of Head-Down Bed-Rest

Alba Martín-Yebra*, Enrico G. Caiani, Violeta Monasterio, Alessandro Pellegrini, Pablo Laguna and Juan Pablo Martínez

Abstract—Reports of ventricular arrhythmias during space-flights raise the question of whether microgravity increases sudden cardiac death risk. We aimed at studying changes in T-wave alternans (TWA) together with Autonomic Nervous System activity during tilt-table (TT) testing before and after five days of -6° head-down bed-rest (HDBR), simulating exposure to microgravity. ECG signals were obtained in 22 males during TT, before and immediately after the end of HDBR, analyzed for multilead TWA detection and spectral analysis of heart rate variability. No differences in TWA indices before and after HDBR were found. However, subjects with lower orthostatic tolerance time after HDBR presented higher TWA values and unbalanced sympato-vagal response to TT already before HDBR.

I. INTRODUCTION

Microgravity leads to cardiovascular deconditioning, inducing significant changes in autonomic and cardiovascular systems, which may adversely influence cardiac repolarization with effects on cardiac rhythm disturbances [1], [2]. In this context, T-wave alternans (TWA) reflecting temporal and spatial repolarization heterogeneity and regarded as a noninvasive risk marker for predicting sudden cardiac death and ventricular vulnerability [3], could be affected. The head-down (6°) bed-rest (HDBR) model represents a way to induce and study the effects of exposure to simulated microgravity on the cardiovascular system.

We hypothesized that simulated microgravity could induce changes in cardiac repolarization and autonomic nervous system (ANS), thus increasing arrhythmia susceptibility and being manifested by an increase in TWA. Accordingly, we studied both changes in TWA and the role of sympathetic drive during tilt-test (TT) before and after 5 days of HDBR.

II. MATERIALS AND METHODS

A. Data acquisition and preprocessing

Twenty-two male healthy subjects (age range 21–43 years) were recruited in the context of the European Space Agency

This work was supported by CIBER-BBN through Instituto de Salud Carlos III and FEDER (Spain), project TEC2010-21703-C03-02 funded by MINECO and FEDER, and by DGA (Aragón) and European Social Fund (EU) through Grupo Consolidado BSICoS. This research has been performed thanks to the contribution of the Italian Space Agency (contract n. I/047/10/0, recipient Dr. EG Caiani). We are extremely grateful to all the personnel of ESA, MEDES and DLR involved in the bed rest studies for the support to the realization of our experiment, as well as to the experimental subjects for their dedicated collaboration.

A. Martín-Yebra, A. Pellegrini and E.G. Caiani are with Dipartimento di Elettronica, Informazione e Bioingegneria, Politecnico di Milano, Italy (*corresponding author e-mail: alpapilar.martin@polimi.it).

A. Martín-Yebra, P. Laguna and J.P. Martínez are with Aragón Institute of Engineering Research, Universidad de Zaragoza and CIBER-BBN, Spain.

V. Monasterio is with the School of Computer Science, San Jorge University, Zaragoza, Spain.

(ESA) bed-rest studies. The experiment included 5 days of pre-bed-rest hospitalization (PRE), 5 days of HDBR and 5 days of post-bed-rest recovery (POST).

ECG signals considered in this study were acquired using 12-lead 24-hours Holter digital recorder ($f_s=1000$ Hz, H12+, Mortara Instrument Inc., Milwaukee, WI) during TT, before and at the end of HDBR. TT consisted of 5 min in supine position, followed by head-up tilt (80° , 30 min) with additional lower body negative pressure applied after this period. Stop criteria were defined as very low BP, extreme tachycardia or clinical symptoms.

Preprocessing of ECG recordings included QRS detection using a wavelet-based ECG delineator [4]. Baseline wander was removed in each lead with a cubic spline interpolation technique. ECGs were low-pass filtered ($f_c=15$ Hz) and down-sampled to remove noise out of TWA frequency range.

B. TWA analysis

Automatic TWA analysis was performed using a multilead scheme based on Periodic Component Analysis (π CA) to find the optimal linear transformation from the 8 standard leads (V1-V6, I and II) to a new set of leads (T1 ... T8) where the 2-beat periodicity of the ST-T complex was maximized. TWA, if present, is mostly projected into the first lead T1. Laplacian Likelihood Ratio (LLR) method for TWA estimation was then applied to the combined lead [5].

ECG records were analyzed in segments of 32 beats with a 50% overlap. The TWA amplitude in each segment (V_k) was defined as the absolute value of the mean amplitude of the estimated TWA waveform, computed as the median difference between ST-T complexes of even and odd beats. A stability criterion in HR was also imposed and only suitable segments were selected for automatic analysis. Three average TWA amplitude indices were computed: 1) TWA_{BAS} : during the 4 minutes preceding the TT (baseline); 2) TWA_{TILT} : during the first 4 minutes of TT; 3) TWA_{REC} : during the first 4 minutes once supine position was restored.

C. Heart rate variability

Heart rate (HR) signal, derived from QRS detection marks, was corrected by integral pulse frequency modulation model [6] and sampled at 4 Hz, obtaining the new signal $d_{HR}(n)$. Heart rate variability (HRV) was defined as $d_{HRV}(n) = d_{HR}(n) - d_{HRM}(n)$, where $d_{HRM}(n)$ is the mean heart rate estimated by low-pass filtering $d_{HR}(n)$ with a cut-off frequency of 0.03 Hz. Three segments were considered to avoid HR transitions: 1) *BAS*, from minute 4 up to 30 sec before the tilt; 2)

TABLE I
TWA AND HRV INDICES COMPUTED BEFORE (PRE) AND AFTER 5 DAYS
OF HDBR (POST). RESULTS ARE EXPRESSED AS MEDIAN
(INTER-QUARTILE RANGE)

	PRE-HDBR	POST-HDBR
HR_{BAS} (beats/min)	61.0 (18.3)	67.7 (21.3) ‡
TWA_{BAS} (μV)	7.74 (8.3)	5.6 (5.4)
LF_{BAS} (s^2)	0.0011 (0.0005)	0.0008 (0.0006) †
HF_{BAS} (s^2)	0.00046 (0.00047)	0.00026 (0.00016) ‡
LFn_{BAS}	0.70 (0.19)	0.76 (0.13) †
LF/HF_{BAS}	2.38 (2.66)	3.17 (2.18) †
HR_{TILT} (beats/min)	85.1 (14.8)	98.5 (21.6) ‡
TWA_{TILT} (μV)	5.6 (4.6)	3.2 (3.9)
LF_{TILT} (s^2)	0.0018 (0.0014)	0.0008 (0.0011) †
HF_{TILT} (s^2)	0.00027 (0.00028)	0.00013 (0.00015) ‡
LFn_{TILT}	0.85 (0.13)	0.87 (0.09)
LF/HF_{TILT}	5.62 (10.20)	6.84 (8.01)
HR_{REC} (beats/min)	78.7 (14.3)	80.9 (19.9)
TWA_{REC} (μV)	5.6 (8.4)	6.9 (4.8)
LF_{REC} (s^2)	0.0012 (0.0021)	0.0015 (0.0010)
HF_{REC} (s^2)	0.00056 (0.0014)	0.00036 (0.00036)
LFn_{REC}	0.71 (0.20)	0.78 (0.15) †
LF/HF_{REC}	2.43 (2.60)	3.58 (3.15) †

‡: $p < 0.005$, †: $p < 0.05$, Wilcoxon signed-rank test, PRE vs POST

TILT, from 30 sec after TT starts to 4 min; 3) *REC*, from 30 sec after supine position was restored to minute 4 of recovery. For each segment, the power spectrum of $d_{HRV}(n)$ was computed by using the periodogram estimator. The power in the LF (0.04 to 0.15 Hz) and HF (0.15 to 0.4 Hz) bands, P_{LF} and P_{HF} , was estimated by integrating the power spectrum in the corresponding bands. Then, P_{LF} was expressed also in normalized units (P_{LFn}) and the ratio of sympatho-vagal absolute power ($R_{LF/HF}$) was computed.

III. RESULTS

In Table I, results of HR, TWA indices and HRV analysis are presented. An increase in HR was found at *BAS* and *TILT* when comparing PRE vs POST. Despite this increment, and although it is known that TWA is a HR-dependent phenomenon, TWA indices were unchanged. As regards sympathetic activity, during *BAS* and *REC*, an increment in P_{LFn} and $R_{LF/HF}$ was found, evidencing a shift in the ANS balance at POST in these two phases but not during *TILT*.

Significant differences between PRE and POST in orthostatic tolerance times (OTT) were found (median (IQR): 38.7 (6.5) vs 8.3 (26.8) min, Wilcoxon signed-rank test, $p < 0.001$), but with no correlation with TWA indices TWA_{TILT} and TWA_{REC} .

Based on OTT at POST, we subdivided the subjects into two groups: 1) *GShort*, with $OTT \leq 30$ min; 2) *GLong*, with $OTT \geq 30$ min. Three subjects were excluded as they presented short OTT (7.9, 22 and 24.3 min) already at PRE and even shorter OTT (3.5, 3.9, 8.3 min respectively) at post. Thus, 12 subjects were included in the *GShort* group, whereas the remaining 7 were included in *GLong*.

When comparing these groups, TWA_{TILT} at POST was higher in *GShort* than in *GLong* (4.6 (7.02) vs. 1.92 (2.12) μV , Mann-Whitney test, $p = 0.028$), which may suggest that subjects who showed higher orthostatic intolerance to tilt induced by HDBR presented a higher TWA index (or higher electrical instability in the repolarization phase). No other differences were found in terms of TWA, while at PRE during *TILT* both P_{LFn} and

$R_{LF/HF}$ increased compared to *BAS* in *GShort* (0.68 (0.17) vs 0.91 (0.09) and 2.13 (2.44) vs 10.97 (10.33), Wilcoxon signed-rank test, $p = 0.002$, respectively) but not in *GLong* (0.74 (0.19) vs 0.80 (0.17) and 2.85 (2.52) vs 4.04 (5.91) respectively), suggesting that sympathetic drive during TT increased in particular in *GShort*. After HDBR, ANS response to tilt was still evident in both groups, but without differences among them.

IV. DISCUSSION AND CONCLUSION

We studied the potential changes in ventricular repolarization induced by HDBR by means of TWA and its possible relationship with ANS activity under TT orthostatic stress conditions. To our knowledge, this is the first study that attempts a description of TWA phenomenon during tilt test.

A significant shortening in OTT after HDBR was found, indicative of initial alterations in ANS activity after simulated microgravity. Based on these OTT changes and on the proposed subject classification, subjects classified as *GShort* presented a significant increase in both P_{LFn} and $R_{LF/HF}$ at PRE in contrast to the *GLong* group. These results are in agreement with previous studies on syncopal and postural orthostatic tachycardia syndrome patients, in which the reaction of sympathetic tone to orthostasis was more severe compared to control subjects [7]. In addition, higher TWA_{TILT} values were found in this group, suggesting higher electrical instability induced by HDBR associated to cardiac deconditioning.

Despite these differences, and the expected fact that higher TWA values are associated to higher sympathetic activation, TWA_{TILT} was lower than TWA_{BAS} , while P_{LFn} and $R_{LF/HF}$ always increased with tilt. Nonetheless, the relationship between both phenomena remains controversial [8], [9], and further research would be needed to further elucidate it.

Higher TWA values during tilt were observed in subjects with shorter OTT, which may be an indication of initial changes in the myocardial substrate. A 5-day period, however, may not be long enough to increase ventricular repolarization's heterogeneity up to the point of altering TWA parameters.

REFERENCES

- [1] Charles JB JB, Bungo MW, Fortner W. "Cardiopulmonary function. Space. Physiology and Medicine", 3rd Edition 1994; 286-304
- [2] Hawkins WR, Zieglschmid JF. "Clinical aspects in crew health. Biomedical results of Apollo 1975"; (NASA SP-386) 286-304
- [3] Verrier RL et al. "Microvolt T-wave alternans physiological basis, methods of measurement, and clinical utility - consensus guideline by International Society for Holter and Noninvasive Electrocardiology". *J Am Coll Cardiol* 2011; 58(13):1309-24
- [4] Martínez JP, Almeida R, Olmos S, Rocha AP, Laguna P. "Wavelet-based ECG delineator: evaluation on standard databases". *IEEE Trans Biomed Eng* 2004;51:570-581
- [5] Monasterio V, Clifford GD, Laguna P, Martínez JP. "A multilead scheme based on periodic component analysis for T-wave alternans analysis in the ECG". *Ann Biomed Eng* 2010;38(8):2532-2541
- [6] Mateo J, Laguna P. "Analysis of heart rate variability in the presence of ectopic beats using the heart timing signal". *IEEE Trans Biomed Eng* 2003;50:334-343
- [7] Duplyakov D, Golovina G, Sysuenkova E, Garkina S. "Can the result of a tilt test be predicted in the first five minutes?" *Cardiology Journal*. 2011;18(5):521-6
- [8] Schwartz PJ, Malliani A. "Electrical alternation of the T-wave: clinical and experimental evidence of its relationship with the sympathetic nervous system and with the long QT syndrome". *Am Heart J* 1975;89:45-50
- [9] Kaufman ES, Mackall JA, Julka B, Drabek C, Rosenbaum, DS. "Influence of heart rate and sympathetic stimulation on arrhythmogenic TWA". *Am J Physiol Heart Circ Physiol* 2000; 279: 1248-1255